

Some parents are concerned about ingredients contained in vaccines, specifically aluminum, mercury, gelatin and antibiotics. However, parents can be reassured that ingredients in vaccines are minuscule and necessary.

Q. Why is aluminum in vaccines?

A. Aluminum is used in vaccines as an *adjuvant*. Until recently, it was the only class of adjuvants approved for use in the United States. In 2009, a second adjuvant, known as monophosphoryl lipid A, was also approved for use in the United States. Adjuvants enhance the immune response by allowing for lesser quantities of active ingredients and, in some cases, fewer doses.

Aluminum

Aluminum salts have been used as adjuvants in vaccines in the United States since the 1930s. Some people wonder whether aluminum in vaccines is harmful — the facts are reassuring.

First, aluminum is present in our environment; the air we breathe, the water we drink and the food we eat all contain aluminum.

Second, the quantity of aluminum in vaccines is small. For example, in the first six months of life, babies receive about 4 milligrams* of aluminum if they get all of the recommended vaccines. However, during this same period they will ingest about 10 milligrams of aluminum if they are breastfed, 40 milligrams if they are fed regular infant formula, and up to 120 milligrams if they are fed soy-based infant formula.

Some people wonder about the difference between aluminum injected in vaccines versus aluminum ingested in food. Typically, infants have between one and five nanograms (billionths of a gram) of aluminum in each milliliter of blood. Researchers have shown that after vaccines are injected, the quantity of aluminum detectable in an infant's blood does not change and that about half of the aluminum from vaccines is eliminated from the body within one day. In fact, aluminum causes harm only when kidneys are not functioning properly or at all (so aluminum cannot be effectively eliminated) AND large quantities of aluminum, such as those in antacids, are administered.

Monophosphoryl lipid A

Monophosphoryl lipid A was isolated from the surface of bacteria and detoxified, so that it cannot cause harm. This adjuvant has been tested for safety in tens of thousands of people and is currently used in one of the HPV vaccines (i.e., Cervarix).

*A milligram is one-thousandth of a gram and a gram is the weight of one-fifth of a teaspoon of water.

Q. Why is formaldehyde in vaccines?

A. Formaldehyde is a by-product of vaccine production. Formaldehyde is used during the manufacture of some vaccines to inactivate viruses (like polio and hepatitis A viruses) or bacterial toxins (like diphtheria and tetanus toxins). While most formaldehyde is purified away, small quantities remain.

Because formaldehyde is associated with the preservation of dead bodies, its presence in vaccines seems inappropriate. However, it is important to realize that formaldehyde is also a by-product of protein and DNA synthesis, so it is commonly found in the bloodstream. The quantity of formaldehyde found in blood is 10 times greater than that found in any vaccine.

Q. Why is gelatin in vaccines?

A. Gelatin is used in some vaccines as a *stabilizer*. Stabilizers are added to vaccines to protect the active ingredients from degrading during manufacture, transport and storage. Gelatin, which is made from the skin or hooves of pigs, is concerning because some people (about 1 of every 2 million) might have a severe allergic reaction to it.

Also, because religious groups, such as Jews, Muslims and Seventh Day Adventists follow dietary rules that prohibit pig products, some parents are concerned about using vaccines that contain gelatin. However, all religious groups have approved the use of gelatin-containing vaccines for their followers for several reasons: First, vaccines are injected, not ingested (except the rotavirus vaccine, which does not contain gelatin). Second, gelatin in vaccines has been highly purified and hydrolyzed (broken down by water), so that it is much smaller than that found in nature; therefore, religious leaders believe it to be different enough that it does not break the religious dietary laws. Finally, leaders from these religious groups believe that the benefits of receiving vaccines outweigh adherence to religious dietary laws.

Q. What about the cumulative effect of vaccine ingredients when my child receives multiple vaccines in a single day?

A. Questions about the cumulative effect when multiple vaccines are given on the same day are reasonable. However, several sources of information provide reassurance:

- A study by Michael Smith and Charles Woods showed that 7- to 10-year-old children who received vaccines according to the recommended schedule as infants did not have neuropsychological delays, such as speech and language delays, verbal memory, fine motor coordination, motor or phonic tics, and intellectual functioning.
- If a new vaccine is added to the schedule at a time when other vaccines are given, studies must be completed to show that neither vaccine interferes with the safety or ability of the other to work. Known as concomitant use studies, these studies are numerous and extensive, offering additional information regarding interference of vaccine ingredients or effects caused by too much of an ingredient.
- Studies of the immune system estimate that we can respond to about 10,000 different immunologic components at any one time; the number of immunologic components contained in all of the vaccines recommended for young children today is less than 200 immunologic components.
- Finally, additives in vaccines, such as aluminum, have been studied regarding how they are processed in the body as well as what levels are toxic. For example, people who suffer toxic effects of aluminum must have had long-term exposure to aluminum (months or years) as well as non-functioning or improperly functioning kidneys.

With all of this information, we can conclude that multiple vaccines given in one day are not overwhelming an infant's immune system.

Vaccine Ingredients: What you should know

Q. Why is mercury in vaccines?

A. Mercury is contained in some multi-dose preparations of influenza vaccine as a *preservative*. Preservatives prevent contamination with bacteria. Early in the 20th century, most vaccines were packaged in vials that contained multiple doses. Doctors and nurses would draw up a single dose and place the vaccine back in the refrigerator. Unfortunately, sometimes bacteria would inadvertently enter the vial and cause abscesses at the site of injection or bloodstream infections that were occasionally fatal. Preservatives, originally added in the 1930s, solved this problem.

The most common preservative used was thimerosal, a mercury-containing compound. As more vaccines were given, children received greater quantities of thimerosal. By the late 1990s, the American Academy of Pediatrics and the Public Health Service requested that mercury be removed from vaccines to make “safe vaccines safer.” No evidence existed to suggest that thimerosal was causing harm, but they wanted to be cautious. Unfortunately, their caution worried parents who wondered whether mercury in vaccines was causing subtle signs of mercury poisoning or autism. Addressing these concerns, scientists performed several studies, all of which showed that thimerosal at the level contained in vaccines hadn’t caused harm.

Further, because mercury is a naturally occurring element found in the earth’s crust, air, soil and water, we are all exposed to it. In fact, infants who are exclusively breastfed ingest more than twice the quantity of mercury than was contained in vaccines. Today, breastfed infants ingest 15 times more mercury in breast milk than is contained in the influenza vaccine.

Q. Are some vaccines made using fetal cells?

A. Fetal cells are used to make five vaccines: rubella, chickenpox, hepatitis A, shingles and rabies. Fetal cells used to grow the vaccine viruses were isolated from two elective abortions performed in Sweden and England in the early 1960s. Further abortions are not necessary as the cells isolated in the 1960s continue to be maintained in laboratory cultures.

Some parents wonder why scientists would choose to use fetal cells at all. There are several reasons for this. First, viruses, unlike bacteria, require cells to grow, and human cells are often better than animal cells at supporting the growth of human viruses. Second, fetal cells are different from other types of cells in that they are virtually immortal, meaning they can reproduce many, many times before dying. Other cells reproduce only a limited number of times before they die.

Some questions have been raised regarding the use of vaccines grown in fetal cells by people whose religious beliefs are against abortions. In 2005, when Pope Benedict XVI was head of the Catholic Church’s Congregation of the Doctrine of Faith, this question was addressed; it was determined that parents who chose not to give vaccines derived from these cells would be in “more proximate cooperation with evil” than those who gave their children the vaccines in question because of the life-saving nature of vaccines. Similarly, the National Catholic Bioethics Center determined that use of vaccines grown in fetal cells isolated from historic abortions was morally acceptable.



This information is provided by the Vaccine Education Center at The Children’s Hospital of Philadelphia. The Center is an educational resource for parents and healthcare professionals and is composed of scientists, physicians, mothers and fathers who are devoted to the study and prevention of infectious diseases. The Vaccine Education Center is funded by endowed chairs from The Children’s Hospital of Philadelphia. The Center does not receive support from pharmaceutical companies.

Q. Do ingredients in vaccines cause allergic reactions?

A. In addition to gelatin, other ingredients in vaccines such as egg proteins, antibiotics and yeast proteins might cause an allergic reaction. Latex used in vaccine packaging is also a concern related to allergies.

Egg proteins

Because the influenza and yellow fever vaccines are grown in eggs, the final products may contain egg proteins. Advances in protein chemistry have resulted in significantly lower quantities of egg proteins in the influenza vaccine; therefore, people with egg allergies can now get influenza vaccine. However, it is recommended that egg-allergic vaccine recipients remain in the office for 30 minutes after getting the influenza vaccine in case of any reaction.

Antibiotics

Antibiotics are used to prevent bacterial contamination during production of some vaccines. However, the types of antibiotics used in vaccines, such as neomycin, streptomycin, polymyxin B, chlortetracycline and amphotericin B, are not those to which people are usually allergic.

Yeast proteins

A couple of viral vaccines are made in yeast cells; these include hepatitis B vaccine and one of the human papillomavirus vaccines (i.e., Gardasil). Although the vaccine is purified away from the yeast cells, about 1 to 5 millionths of a gram remain in the final product. The good news is that people who are allergic to bread or bread products are not allergic to yeast, so the risk of allergy from yeast is theoretical.

Latex packaging

A small number of vaccines are packaged with materials that include latex. While it is rare that patients have a reaction to latex in vaccine packaging, people with latex allergies should consult with their allergy doctor before getting any vaccines packaged in this way.

Selected References

Aluminum: Baylor NW, Egan W, Richman P. Aluminum salts in vaccines — U.S. perspective. *Vaccine*. 2002;20:S18-S23.

Formaldehyde: Epidemiology of chronic occupational exposure to formaldehyde: report of the ad hoc panel on health aspects of formaldehyde. *Toxicology and Industrial Health*. 1988;4:77-90.

Gelatin: Atkinson WL, Kroger AL, and Pickering LK. General Immunization Practices. In: Plotkin SA, Orenstein WA, and Offit PA, eds., *Vaccines Sixth Edition*. Saunders Elsevier, 2012.

Cumulative effects: Smith MJ and Woods CR. On-time vaccine receipt in the first year does not adversely affect neuropsychological outcomes. *Pediatrics*. 2010;125(6):1134-1141.

Thimerosal: Gerber, JS and Offit, PA. Vaccines and autism: A tale of shifting hypotheses. *Clinical Infectious Diseases*. 2009;48:456-461.

Fetal cells: Offit PA. *Vaccinated: One man’s quest to defeat the world’s deadliest diseases*. New York: Harper Perennial, 2007.

Allergic reactions: Offit PA, Jew RK. Addressing parents’ concerns: do vaccines contain harmful preservatives, adjuvants, additives, or residuals? *Pediatrics*. 2003;112:1394-1401.

Multiple Vaccines: Offit PA, Quarles J, Gerber MA, Hackett CJ, Marcuse EK, Kollman TR, Gellin BG, Landry S. Addressing parents’ concerns: Do multiple vaccines overwhelm or weaken the infant’s immune system? *Pediatrics*. 2002 Jan;109(1):124-129.

 The Children’s Hospital
of Philadelphia®

 VACCINE EDUCATION CENTER

vaccine.chop.edu

The Children’s Hospital of Philadelphia, the nation’s first pediatric hospital, is a world leader in patient care, pioneering research, education and advocacy.